

10/019993

531 Rec'd PCT 09 NOV 2001

Express Mail No. ET401306226US

AROMATIC AMIDES

FIELD OF THE INVENTION

This invention provides aromatic amides which are useful CNS agents, especially for treating depression, pain, anxiety, schizophrenia and seizure disorders.

BACKGROUND OF THE INVENTION

Disorders of the central nervous system have become one of the most common and most debilitating diseases currently afflicting mankind. Specific disorders such as depression and schizophrenia are now known to be common afflictions, and are routinely diagnosed. These diseases result in significant losses of an individual's ability to work and to carry out normal daily activities, and in many cases require long term hospitalization or institutionalization. Only recently have new treatments, such as the selective serotonin re-uptake inhibitors for example, become available and are effective for many people. Unfortunately, such agents are not effective for all cases of depression, and indeed can lead to significant adverse reactions in some patients.

Other CNS disorders, such as chronic pain and seizure disorders, are only marginally treatable, and such treatments often are associated with unacceptably high health risks, for instance long term use of narcotic analgesics to treat chronic pain generally results in addiction to the drug being employed, the results of which can be devastating to the patient.

Accordingly, the need continues for new medicines that will effectively treat CNS disorders without imposing unacceptable liability and risk issues. I have now discovered a series of aromatic amides which can be utilized to treat these CNS disorders, and which have a very good risk-to-benefit ratio. The invention compounds are alkyl amides having an aromatic group attached to the amide nitrogen atom.

Several N-aryl alkylamides are known in the prior art. For example, Ronsisvalle *et al.* described a series of analgesic N-thienyl acetamides in Eur. J. Med. Chem. 3: 553-559, 1998.

US Patent No. 4,203,988 discloses certain N-pyridyl amide derivatives as inhibitors of gastric secretion, while US No. 3,163,645 discloses N-pyridyl amides as analgesics. US No. 5,372,931 discloses N-alkoxyphenyl and N-alkoxynaphthyl amides as useful in certain analytical and diagnostic methods.


- 5 Elslager *et al.*, in J. Med. Chem. 9: 378-91, 1966, describe certain N-naphthyl amides as useful as intermediates in the synthesis of arylazo substituted naphthyl alkylenediamines. Similarly, Elslager *et al.*, described certain N-quinolyl amides in J. Med. Chem. 12: 600-7, 1966.

10 The compounds provided by this invention are characterized as novel N-aryl amides having good CNS activities, and are thus useful for treating depression, anxiety, pain, schizophrenia, and seizure disorders such as epilepsy.

SUMMARY OF THE INVENTION

This invention provides N-aryl alkylamides defined by Formula I


15





wherein :

- R^1 is hydrogen, C_1 - C_4 alkyl, or C_2 - C_4 alkenyl;
- 20 R^2 and R^3 independently are hydrogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, phenyl or benzyl, or taken together with the nitrogen to which they are attached complete a ring having from 4 to 7 ring atoms, one optionally being oxygen; X is $(\text{CH}_2)_n$, $\text{CHMe}-(\text{CH}_2)_{n-1}$ or $(\text{CH}_2)_{n-1}-\text{CHMe}$, n is 1, 2 or 3;
- R^4 is an aromatic or heteroaromatic group selected from

25



wherein R^5 is hydrogen, halogen, C_1 - C_4 alkyl, nitro, N_3 or CF_3 and R^6 is hydrogen, C_1 - C_4

5 alkyl, $-(C=O)Me$, $-(C=O)NH_2$,  or , and the pharmaceutically acceptable salts thereof.

Preferred invention compounds have Formula I wherein R^1 , R^2 and R^3 independently are C_1 - C_4 alkyl, and R^4 is naphthyl, substituted naphthyl, fluorene or substituted fluorene.

10 Also preferred are the compounds of Formula I wherein n is 2 or 3.

Another embodiment of this invention is a pharmaceutical formulation comprising a compound of Formula I admixed with a pharmaceutically acceptable carrier, diluent or carrier therefor.

The compounds of the instant invention are useful for the treatment of CNS
15 disorders including neurodegenerative disorders, pain, depression, convulsions, anxiety, schizophrenia and seizures.

Neurodegenerative disorders include, for example, Alzheimer's disease, Huntington's disease, Parkinson's disease, and Amyotrophic Lateral Sclerosis.

The present invention also covers treating neurodegenerative disorders termed
20 acute brain injury. These include but are not limited to: stroke, head trauma, and asphyxia.

Stroke refers to a cerebral vascular disease and may also be referred to as a cerebral vascular incident (CVA) and includes acute thromboembolic stroke. Stroke includes both focal and global ischemia. Also, included are transient cerebral ischemic attacks and other cerebral vascular problems accompanied by cerebral ischemia. A patient undergoing carotid endarterectomy specifically or other cerebrovascular or vascular surgical procedures in general, or diagnostic vascular procedures including cerebral angiography and the like.

Other incidents are head trauma, spinal cord trauma, or injury from general anoxia, hypoxia, hypoglycemia, hypotension as well as similar injuries seen during procedures from embole, hyperfusion, and hypoxia.

The instant invention would be useful in a range of incidents, for example, during cardiac bypass surgery, in incidents of intracranial hemorrhage, in perinatal asphyxia, in cardiac arrest, and status epilepticus.

Pain refers to acute as well as chronic pain.

Acute pain is usually short-lived and is associated with hyperactivity of the sympathetic nervous system. Examples are postoperative pain and allodynia.

Chronic pain is usually defined as pain persisting from 3 to 6 months and includes somatogenic pains and psychogenic pains. Other pain is nociceptive.

Still other pain is caused by injury or infection of peripheral sensory nerves. It includes, but is not limited to pain from peripheral nerve trauma, herpes virus infection, diabetes mellitus, causalgia, plexus avulsion, neuroma, limb amputation, and vasculitis. Neuropathic pain is also caused by nerve damage from chronic alcoholism, human immunodeficiency virus infection, hypothyroidism, uremia, or vitamin deficiencies. Neuropathic pain includes, but is not limited to pain caused by nerve injury such as, for example, the pain diabetics suffer from.

Psychogenic pain is that which occurs without an organic origin such as low back pain, atypical facial pain, and chronic headache.

Other types of pain are: inflammatory pain, osteoarthritic pain, trigeminal neuralgia, cancer pain, diabetic neuropathy, restless leg syndrome, acute herpetic and postherpetic neuralgia, causalgia, brachial plexus avulsion, occipital neuralgia, gout, phantom limb, burn, IBS and other forms of neuralgia, neuropathic and idiopathic pain syndrome.


The compounds of the invention are also useful in the treatment of depression. Depression can be the result of organic disease, secondary to stress associated with personal loss, or idiopathic in origin. There is a strong tendency for familial occurrence of some forms of depression suggesting a mechanistic cause for at least some forms of depression. The diagnosis of depression is made primarily by quantification of alterations in patients' mood. These evaluations of mood are generally performed by a physician or quantified by a neuropsychologist using validated rating scales, such as the Hamilton Depression Rating Scale or the Brief Psychiatric Rating Scale. Numerous other scales have been developed to quantify and measure the degree of mood alterations in patients with depression, such as insomnia, difficulty with concentration, lack of energy, feelings of worthlessness, and guilt. The standards for diagnosis of depression as well as all psychiatric diagnoses are collected in the Diagnostic and Statistical Manual of Mental Disorders (Fourth Edition) referred to as the DSM-IV-R manual published by the American Psychiatric Association, 1994.

The invention also provides a method for treating CNS disorders in mammals, comprising administering a CNS effective amount of a compound of Formula I to a mammal in need of treatment.


As used herein, the term "C₁-C₄ alkyl" means straight and branched carbon chains having from 1 to 4 carbon atoms, for example methyl, ethyl, n-propyl, isopropyl, *n*-butyl, *sec*-butyl and *tert*-butyl.

30 "Halo" means fluoro, chloro, bromo and iodo.


"Substituted aryl" and "substituted heteroaryl" means any of the cyclic ring systems described above having R^5 other than hydrogen, for example where R^5 is halo, C_1 - C_4 alkyl, nitro or CF_3 . Typical substituted aryl and substituted heteroaryl groups thus include 3-chloronaphthyl, 4-nitronaphthyl, 4-nitrobenzofuranyl, 3-methylbenzothieryl, and 1-methyl-3-trifluoromethyl indole. These are compounds of Formula I wherein R^4 is a cyclic, bicyclic or tricyclic aromatic or heteroaromatic group bearing a substituent defined as R^5 , where R^5 is other than hydrogen. The group



is a naphthyl ring which can be attached to the amide nitrogen (of Formula I) at any ring position. This ring can be substituted at any available ring position by the group R^5 . Specific examples include :




Specific examples of the group:




15


include:




Specific examples of the group:



include:




and




5


Specific examples of the group:




include




Specific examples of the group:




10 include




Specific examples of the group:




include



,




and




5


Specific examples of the group:




include




,




and




Specific examples of the group:




10 include




,




,




,




,



and




Specific examples of the group:




include


5



Specific examples of the group:



10 include




The compounds of this invention are amines and as such they readily form pharmaceutically acceptable salts by reaction with common inorganic and organic acids. Typical acids commonly used to form salts include hydrochloric, nitric, phosphoric, and sulfuric acid, as well as acetic, citric, malonic, tartaric, succinic, salicylic, methanesulfonic, oxalic and benzoic acid. Any common inorganic or organic acid can be utilized to form the pharmaceutically acceptable salts of this invention, and the specific acid to be utilized is well within the skill of the art.

The compounds provided by this invention can be prepared by any of several methods well known to those of ordinary skill in the art of organic chemistry. In a typical synthesis, an N-aryl alkyl diamine is acylated, for example by reaction with an aryl halide, or by


20

coupling an aryl-acid to the amide in the presence of a common peptide coupling reagent such as DCC (dicyclohexylcarbodiimide). Such synthesis can be illustrated by Scheme 1, in which an alkyl diamine is first prepared by reacting a halo substituted acyl halide with an amine HNR^2R^3 , to give the corresponding halo substituted amide, reacting the halo substituted amide with an aryl amine ArNH_2 to give an arylaminoamide, reducing the amide carbonyl to give the corresponding arylamino alkylamine, and then acylating the arylamino nitrogen atom to give a compound of Formula II. The synthetic sequence is illustrated in scheme 1:



10

An alternative method for preparing the invention compounds comprises alkylating a terminal primary or secondary amine of the formula




15

where one or both of R^2 and R^3 are hydrogen, by reaction with an alkylating agent such as an alkyl halide. The reaction is depicted by scheme 2, which illustrates the synthesis of the primary or secondary amine according to the general scheme shown above, followed by a reaction with a common alkylating agent.

20

Scheme 2



- 5 In the above scheme, the halo substituted acid halide is reacted with an amine bearing a group that is easily removed, such as benzyl. This is a normal acylation reaction that is typically carried out in a solvent such as dichloromethane or toluene, and generally is complete within 30 min to 1 h when carried out at a temperature of about 30°C to about 60 °C. The resulting amide is readily isolated by removing the solvent, and is subsequently
- 10 reacted with an amine R^4NH_2 in the presence of a base such as sodium carbonate or triethylamine, and typically in a solvent such as N,N-dimethylformamide or diethyl ether. The resulting amino substituted amide is readily isolated by removing the solvent, and further purification generally is not required. The amino substituted amide is readily reduced by reaction with a reducing agent such as lithium aluminium hydride or sodium borohydride, thus affording an alkylene diamine. The alkylene diamine is coupled to an
- 15 acyl group, for example by common acylation with an acid anhydride or acid halide (e.g. $\text{R}^1-\text{C}(=\text{O})-\text{O}-\text{C}(=\text{O})-\text{R}^1$ or $\text{R}^1-\text{C}(=\text{O})-\text{halo}$, or by reacting the free acid R^1COOH with the amine using a coupling reagent such as dicyclohexylcarbodiimide (DCC).
- The corresponding amide is next converted to a primary or secondary amine, for instance
- 20 by removing a group such as benzyl by normal catalytic hydrogenation. The resulting amine is reacted with a common alkylating agent such as an alkyl halide (R^3-halo) and the


resulting product of Formula I is isolated by removing any reaction solvent and excess alkylating agent. The invention compound can be further purified if desired by routine methods such as crystallization, for example from solvents such as methanol, diethylether, ethyl acetate and the like, or chromatography over solid supports such as silica gel.

5

Still another way to prepare the invention compounds is to start with an aryl amine (R^4NH_2), acylate it with an acyl halide or anhydride to form an amide, and then alkylate the amide with an amino substituted alkyl halide. This process is depicted in Scheme 3 below :

10

Scheme 3



15


These reactions are carried out under normal organic synthetic conditions. For example, an aryl amine such as 3-naphthylamine can be reacted with acetyl chloride in a solvent such as toluene. A base such as triethylamine can be utilized as an acid scavenger if desired. The reaction is substantially complete within 1 to 2 h when carried out at about 30 to 60 °C, and the product amide is readily isolated by removing the reaction solvent. The amine is then alkylated by reaction with an amino substituted amino alkyl halide to produce the invention compound of Formula I.

20

The synthesis of specific invention compounds is further illustrated by the following detailed example. The examples are representative only, and are not intended to limit the invention in any respect.

25

EXAMPLE 1



5 Reagents : (i) propionyl chloride, Et_3N ; (ii) NaH , $\text{Et}_2\text{NCH}_2\text{CH}_2\text{Cl}\cdot\text{HCl}$

N-Propionyl 1-amino-4-chloronaphthalene.

To a stirred solution of 1-amino-4-chloronaphthalene (0.70 g, 3.9 mmol) in dichloromethane (50 ml) was added triethylamine (1.0 ml, 7 mmol), followed by propionyl chloride (0.5 ml, 5.8 mmol). After 20 min the mixture was diluted with ethyl acetate (150 ml) and washed with 2N HCl (100 ml) followed by saturated sodium carbonate (100 ml). The organic phase was separated, dried (MgSO_4) and the solvent removed *in vacuo*. The residue was triturated with a mixture of ethyl acetate and heptane, 130 ml, 3:10) to give 0.62 g (67 %) of the desired compound as a white solid.

15 ^1H NMR 400 MHz (CDCl_3) : τ^m 1.33 (3H, t, $J = 6\text{Hz}$) ; 2.56 (2H, q, $J = 6\text{Hz}$) ; 7.47 (1H, br s) ; 7.52-7.70, 4H, m) ; 7.84 (1H, m) ; 8.32 (1H, m).

MS ES^+ : m/z 236 ($[\text{MH}]^+$, 16%), 234 ($[\text{MH}]^+$, 48%).

IR (thin film) λ_{max} (cm^{-1}) : 1652, 2922, 3300.

20 N-Propionyl, N-(2-diethylaminoethyl)-1-amino-4-chloronaphthalene.

To a stirred solution of N-propionyl 1-amino-4-chloronaphthalene (400 mg, 1.7 mmol) in dry dimethylformamide (40 ml) was added sodium hydride (60% dispersion in oil, 0.2 g, 5 mmol). After 20 min, 2-diethylaminoethylchloride hydrochloride (0.4 g, 2.8 mmol) was added and the mixture stirred for a further 2 h. Water (200 ml) was added and the mixture extracted with ethyl acetate (2 x 100 ml). The organic extracts were combined, dried

25

(MgSO₄) and the solvent removed *in vacuo*. The residue was purified by reverse phase chromatography (methanol:water 7:3) to give 0.27 g (47%) of the desired product as a colorless oil.

¹H NMR 400 MHz (CDCl₃): τ 0.97 (9H, m); 1.80 (1H, m); 2.01 (1H, m); 2.50 (4H, m); 2.69 (2H, t, J = 7Hz); 3.34 (1H, m); 4.33 (1H, m); 7.36 (1H, d, J = 8 Hz); 7.55-7.70 (3H, m); 7.84 (1H, m); 8.34 (1H, d, J = 8 Hz).


MS CI: m/z 233 ([MH]⁺, 100 %).

IR (thin film) λ_{\max} (cm⁻¹): 1667, 2970.

10 Microanalysis for C₁₉H₂₅N₂OCl

Calculated	C	68.56%	H	7.57%	N	8.42%
Found		68.29%		7.78%		8.20%

EXAMPLE 2



15

Reagents : (i) propionyl chloride, Et₃N ; (ii) NaH, Et₂NCH₂CH₂Cl.HCl

N-Propionyl 4-amino-9-fluorenone.

20 To a stirred solution of 4-amino-9-fluorenone (0.20 g, 1.0 mmol) in dichloromethane (40 ml) was added triethylamine (0.5 ml, 3.5 mmol), followed by propionyl chloride (0.5 ml, 5.8 mmol). After 20 min the mixture was diluted with ethyl acetate (150 ml) and washed

with 2N HCl (100 ml) followed by saturated sodium carbonate (100 ml). The organic phase was separated, dried (MgSO₄) and the solvent removed *in vacuo*. The residue was purified by flash chromatography (silica, heptane:ethyl acetate 7:3) to give 164 mg (63%) of the desired material as a yellow oil.

¹H NMR 400 MHz (CDCl₃): τ 1.36 (3H, br t); 2.56 (2H, br q); 7.18-7.38 (4H, m); 7.41-7.60, (2H, m); 7.71 (1H, d, J = 8 Hz); 7.83 (1H, br s).

IR (thin film) ν_{\max} (cm⁻¹): 1659, 1716, 3258.

10 N-Propionyl, N-(2-diethylaminoethyl)-4-amino-9-fluorenone.

N-propionyl 4-amino-9-fluorenone (158 mg, 0.6 mmol) was dissolved in dry dimethylformamide (40 ml) and sodium hydride (60% dispersion in oil, 80 mg, 1.2 mmol). After 20 min, 2-diethylaminoethylchloride hydrochloride (250 mg, 1.4 mmol) was added and the mixture was heated to 80°C. After 10 min the mixture was cooled to room temperature and diluted with water (20 ml). The mixture was diluted with saturated sodium

carbonate (150 ml) and the mixture extracted with ethyl acetate (2 x 70 ml). The organic extracts were combined, dried (MgSO₄) and the solvent removed *in vacuo*. The residue was purified by flash chromatography (silica, dichloromethane:diethyl ether 9:1, and then 1:4) to give 0.16 g (73%) of the desired product as a colorless oil.

¹H NMR 400 MHz (CDCl₃): τ 0.95 (6H, t, J = 7 Hz); 1.05 (3H, t, J = 7 Hz); 2.08 (2H, m); 2.50 (4H, m); 2.69 (2H, m); 3.34 (1H, m); 4.34 (1H, m); 7.30-7.75 (7H, m).

MS CI : m/z 351 ([MH]⁺, 100 %).

IR (thin film) ν_{\max} (cm⁻¹): 1652, 1716, 2970.

25 Microanalysis for C₂₂H₂₆N₂O₂

Calculated	C	75.40%	H	7.48%	N	7.99%
Found		75.55%		7.57%		7.94%

EXAMPLES 3-15

By following the general procedure of Examples 1 and 2, several additional compounds of Formula I were prepared and are described in Table I below.





The compounds of Formula I have been evaluated in standard in vivo and in vitro assays routinely used to measure the ability of test compounds to interact with the central nervous system of animals, thereby establishing their utility for treating CNS disorders such as pain, depression, anxiety and schizophrenia. In a typical assay, compounds are evaluated for their ability to bind to the $\alpha_2\delta$ subunit of the calcium channel found in animal brain tissue. Significant binding to this receptor indicates a compound's analgesic potential.






In another test, compounds were evaluated for their ability to reduce the hyperalgesia effects of carrageenin in the following assay: nociceptive pressure thresholds were measured in the rat paw pressure test using an analgesimeter (Randall L.O. and Selitto J.J., A method for measurement of analgesic activity on inflamed tissue. Arch. Int. Pharmacodyn. 4: 409-419, 1957). Male Sprague-Dawley rats (70-90 g) were trained on this apparatus before the test day. Pressure was gradually applied to the hind paw of each rat. Nociceptive thresholds were determined as the pressure (g) required to elicit paw withdrawal. A cutoff point of 250 g was used to prevent any tissue damage to the paw. On the test day, 2 to 3 baseline measurements were taken before animals were administered 100 μ l of 2 % aqueous carrageenin by intraplantar injection into the right hind paw.




Nociceptive thresholds were taken again 3 h after carrageenin injection to establish that animals were exhibiting hyperalgesia. Animals were orally dosed with a compound of Formula I (by gavage) at 3.5 h after carrageenin injections and nociceptive thresholds were examined at 1 and at 2 h post-carrageenin.




Table 1 presents the biological activity of representative invention compounds when evaluated in the above tests, and in the in vitro $\alpha_2\delta$ binding assay as described by Gee et al. in J. Biol. Chem., 1996; 271: 5776-5879, incorporated herein by reference.

Table 1

			Carrageenin induced thermal hyperalgesia in the rat	
Compound	Structure	IC ₅₀ (μ M) at $\alpha_2\delta$ binding site	%MPE* 1h post dose @30mg/kg p.o.	%MPE* 2h post dose @30mg/kg p.o.
N-Propionyl, N-(2-Diethylaminoethyl)-1-amino-4-chloronaphthalene (Example 1)		0.170	51.5	22.2
N-Propionyl, N-(2-Diethylaminoethyl)-4-amino-9-fluorenone (Example 2)		0.058	1.1	6.4
N-Propionyl, N-(2-Diethylaminoethyl)-1-amino-4-bromonaphthalene (Example 3)		0.065	-2.6	7.7
N-Propionyl, N-(N-Morpholino)-1-amino-4-chloronaphthalene (Example 4)		>10	44.8	30.7

			Carrageenin induced thermal hyperalgesia in the rat	
Compound	Structure	IC ₅₀ (μ M) at $\alpha_2\delta$ binding site	%MPE* 1h post dose @30mg/kg p.o.	%MPE* 2h post dose @30mg/kg p.o.
N-Propionyl, N-(2-(3-diethylamino-propyl))-1-amino-4-chloronaphthalene (Example 5)		5.03	23.3	27.5
N-Propionyl, N-(2-Diethylaminoethyl)-1-amino-4-azidonaphthalene (Example 6)		0.885	N/A	N/A
N-Propionyl, N-(2-Diethylaminoethyl)-3-chlorobenzylamine (Example 7)		1.7	N/A	N/A
N-Propionyl, N-(2-Diethylaminoethyl)-3-bromobenzylamine (Example 8)		4.81	N/A	N/A
N-Propionyl, N-(2-Piperidylethyl)-1-amino-4-chloronaphthalene (Example 9)		> 10	N/A	N/A

			Carrageenin induced thermal hyperalgesia in the rat	
Compound	Structure	IC ₅₀ (μ M) at $\alpha_2\delta$ binding site	%MPE* 1h post dose @30mg/kg p.o.	%MPE* 2h post dose @30mg/kg p.o.
N-Propionyl, N-(2-(3-dimethylamino-propyl))-1-amino-4-chloronaphthalene (Example 10)		2.336	N/A	N/A
N-Propionyl, N-(2-Dimethylaminoethyl)-1-amino-4-chloronaphthalene (Example 11)		5.34	N/A	N/A
N-Propionyl, N-(2-(N-benzyl)-aminoethyl)-1-aminonaphthalene (Example 12)		> 10	29.68	3.13

Compound	Structure	IC ₅₀ (μ M) at $\alpha_2\delta$ binding site	Carrageenin induced thermal hyperalgesia in the rat	
			%MPE* 1h post dose @30mg/kg p.o.	%MPE* 2h post dose @30mg/kg p.o.
N-(2-Diethylamino-ethyl)-N-(7-methyl-quinolin-4-yl)-propionamide (Example 13)		5.47	8.6	1.2
N-Acryloyl, N-(2-Diethylaminoethyl)-1-amino-4-chloronaphthalene (Example 14)		0.177	15.1	0.9
N-Propionyl, N-(2-Diethylaminoethyl)-(1-amino-4-nitronaphthalene (Example 15)		0.800	-5.7	2.0

*MPE: maximum possible effect – set as baseline value prior to treatment with carrageenin

As noted above, the invention compounds of Formula I are typically utilized in the form of pharmaceutical compositions for human therapy of CNS disorders. The compounds can be formulated with any excipient, diluent or carrier commonly utilized in the pharmaceutical art. Such common excipients include potato starch, corn starch, talc, sucrose, lactose, cellulose; flavoring agents such as peppermint, orange flavor and the like. Binders and lubricants such as magnesium stearate, colloidal silicon dioxide and gum tragacanth can be utilized for convenient oral or parenteral administration, for example as tablets, capsules, aqueous solutions, elixirs, syrups, and controlled release patches, pellets and suppositories, as well as solutions for IV, SC and IM injection. The formulations will typically contain from about 5 % to about 95 % of active compound of Formula I (w/w).

The preparations will be administered such that the active ingredient is presented at a dose which is effective to treat a CNS disorder. Such dose will generally be from about 0.1 to about 2000 mg/kg of body weight, typically about 1 mg to about 100 mg/kg. The formulations can be administered from 1 to about 4 times a day, or as otherwise dictated by the particular patient and condition being treated, and the attending medical practitioner.

The compounds of Formula I can additionally be utilized in combination with other active ingredients, for example selective serotonin re-uptake inhibitors such as fluoxetine hydrochloride, and any of the tricyclic antidepressants such as benzazepines and the like.

The following examples further illustrate specific formulations provided by this invention.

EXAMPLE 16

15

Tablets

N-Butyryl, N-(3-dimethylamino-propyl)-5-amino-indole	200 mg
Potato starch	50 mg
Magnesium stearate	25 mg
Talc	25 mg

20

The above ingredients are blended to uniformity and then pressed into a tablet. Such tablets are administered from 1 to 4 times a day to an adult human suffering from depression and in need of treatment.

EXAMPLE 17

25

Capsules

N-pivaloyl 1-amino-2-trifluoromethyl-naphthalene	300 mg
Corn starch	50 mg
Dextrose	50 mg
Magnesium oxide	1 mg

30

The above ingredients are blended to uniformity and filled into an empty telescoping gelatin capsule. Such capsules are administered from 1 to 4 times a day to an adult human suffering from schizophrenia and in need of treatment.

EXAMPLE 18Parenteral solution

	N-propionyl,N-(2-diethylaminoethyl)(1-amino-4-bromonaphthalene),	
5	hydrochloride salt	500 mg
	isotonic saline	qs 1000 ml

The invention compound is dissolved in 1000 ml of isotonic saline and filled into a sterile plastic bottle equipped with a drip tube. The solution is administered IV to a human
10 suffering from chronic pain resulting from colon carcinoma.

EXAMPLE 19Transdermal skin patch

	N-acetyl, N-(3-(N-ethyl-N-isobutyl)aminopropyl-	
15	3-amino-6-bromofluorene	450 mg
	propylene glycol	10 mg
	elastomer	5 mg
	methyl cellulose	50 mg
	sodium carboxymethyl cellulose	25 mg

20

The above ingredients are blended and spread onto an elastic tape. The tape is applied to the skin surface of a mammal to prevent and treat migraine pain.

The compounds of Formula I are useful for treating all conditions resulting from
25 disorders within the central nervous system in animals, including humans. Commonly treated conditions include pain, depression, anxiety and schizophrenia. Other conditions that can be treated according to this invention include seizure disorders, i.e. epilepsy, Huntington's disease, Parkinson's disease, amyotrophic lateral sclerosis (ALS), Alzheimer's disease, migraine, cerebral ischemia, and compulsive disorders such as
30 narcotic addiction, alcoholism, smoking addiction, appetite disorders such as bulimia and obesity, sexual performance, and sleeping disorders.